



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ :

C12N 9/64

A1

(11) International Publication Number:

WO 90/15866

(43) International Publication Date: 27 December 1990 (27.12.90)

(21) International Application Number: PCT/US90/03378

(22) International Filing Date: 13 June 1990 (13.06.90)

(30) Priority data:

365,937

13 June 1989 (13.06.89)

US

(71) Applicant: GENENCOR INTERNATIONAL, INC. [US/
US]; 180 Kimball Way, South San Francisco, CA 94080
(US).(72) Inventors: HEINSOHN, Henry, G. ; 413 Copeland Street,
Pacifica, CA 94044 (US). LORCH, Jeffrey, D. ; 1833
Royal Avenue, San Mateo, CA 94401 (US). HAYENGA,
Kirk, J. ; 1640 Bayridge Way, San Mateo, CA 94402
(US). ARNOLD, Raymond, E. ; 894 Lombard Street,
San Francisco, CA 94133 (US).(74) Agent: SWISS. Gerald, F. ; Burns, Doane, Swecker & Ma-
this, Washington & Prince Streets, P.O. Box 1404, Alex-
andria, VA 22313-1404 (US).(81) Designated States: AT (European patent), AU, BE (Euro-
pean patent), CA, CH (European patent), DE (Euro-
pean patent)*, DK (European patent), ES (European pa-
tent), FI, FR (European patent), GB (European patent),
IT (European patent), JP, LU (European patent), NL
(European patent), NO, SE (European patent).

Published

*With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

Property of
SemBioSys
GENETICS INC.

(54) Title: PROCESSES FOR THE RECOVERY OF NATURALLY PRODUCED CHYMOSIN

(57) Abstract

Disclosed are methods for the recovery and purification of naturally produced chymosin. In particular, disclosed are meth-
ods for the recovery and purification of chymosin from aqueous solutions containing chymosin, pepsin and other contaminants.

PEG/salt - anion 4

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TC	Togo
DK	Denmark			US	United States of America

-1-

5

PROCESSES FOR THE RECOVERY
OF NATURALLY PRODUCED CHYMOSIN

BACKGROUND OF THE INVENTION

1. Field of the Invention.

10 This invention relates to the recovery
and purification of naturally produced
chymosin. In particular, this invention is
directed to methods for the recovery and
purification of chymosin from aqueous
15 solutions containing chymosin which have been
obtained from the extraction of natural
chymosin sources.

2. State of the Art.

20 Chymosin is a known enzyme which is
particularly useful in the preparation of
cheese. While natural sources of chymosin
include calf stomachs, bovine stomachs, goat
stomachs, porcine stomachs, etc., commercial
chymosin has heretofore been primarily
obtained from the fourth stomach of milk fed
25 calves. This is result of the fact that such

-2-

calves produce greater amounts of chymosin as compared to pepsin whereas other chymosin sources generally contain greater amounts of pepsin as compared to chymosin and
5 accordingly, recovery of chymosin from such other sources is more difficult and economically less efficient, i.e., there is less chymosin to recover.

10 However, due to the recent decrease in calf production, the heretofore preferred natural source of chymosin has declined which, in turn, has provided impetus to developing more efficient methods for the recovery and purification of naturally produced chymosin.
15 Specifically, more efficient methods would result in the improved recovery and purification of chymosin from calf stomachs as well as permit the economical recovery and purification of chymosin from other natural
20 chymosin sources.

A major stumbling block to developing such methodology has been the very high level of contaminants found in the chymosin solution obtained from natural chymosin sources. In
25 addition to pepsin, the aqueous extract obtained from these natural chymosin sources contain other contaminants including, for example, other stomach enzymes and proteins. Such contaminants have complicated the
30 development of efficient recovery and purification methodology.

-3-

While numerous methods are disclosed for isolating enzymes from aqueous solutions, such as fermentation beer, none of the references which Applicants are aware of disclose methods for recovering and purifying naturally produced chymosin, especially naturally produced chymosin intermixed with pepsin and other contaminants, which employ a liquid-liquid two phase system.

10 In this regard, U.S. Patent No. 4,144,130 describes the use of (1) a mixture of a high molecular weight unsubstituted or substituted polyalcohol, polyether, polyvinylpyrrolidone or polysaccharide and an inorganic salt, (2) a
15 mixture of at least two of the above high molecular weight polymers to recover intracellular enzymes from an aqueous solution into which they have been released from the cells. When a mixture of polyethylene glycol
20 and an inorganic salt is used, the desired intracellular enzyme goes into the top polyethylene glycol layer while the cell debris and other fermentation products go into the lower salt-containing layer. This
25 reference discloses that the partition coefficients for various enzymes recovered in the glycol layer was about 0.3 when a normal cell mass was treated, which could be increased to about 3 when frozen cells were
30 mixed with water and disintegrated to release their enzymes.

-4-

Similarly, U.S. Patent No. 4,728,613, discloses a process for the recovery of extracellularly produced enzymes, such as protease, amylase and microbial rennet, from whole fermentation beer by using an inorganic salt in combination with a polymer selected from the group consisting of polyethylene glycol, an amine derivative of polyethylene glycol, a carboxylate derivative of polyethylene glycol, polypropylene glycol, an amine derivative of polypropylene glycol, a carboxylate derivative of polypropylene glycol, poly(ethylene glycol) ester, polyethyleneimine, trimethylamino-polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone and mixtures thereof. The examples of this reference disclose achieving partition coefficients of up to about 80 for such extracellular enzymes.

Likewise, Kula et al., "Purification of Enzymes by Liquid-Liquid Extraction", describes numerous methods for the purification of enzymes by liquid-liquid extraction. Among numerous methods disclosed, Kula et al. disclose that the addition of a polyethylene glycol/inorganic salt mixture to an aqueous solution containing the enzyme will form a two phase system wherein the polyethylene glycol phase will contain the enzyme. Kula et al. further disclose at page 111 that the phase forming polymer

-5-

(polyethylene glycol) can be removed from the enzyme by adsorption of the enzyme onto ion exchangers; washing away of the phase forming polymer; and the subsequent recovery of the enzyme.

On the other hand, U.S. Patent No. 4,508,825 discloses that extracellular protease and amylase co-produced during the fermentation of a microorganism capable of producing them are separated by the addition of polyethylene glycol and a cationic epihalohydrin/polyamine copolymer or dextran polymer to the fermentation medium and allowing the polymers to phase separate to form a protease rich phase and an amylase rich phase.

Also, U.S. Patent No. 4,591,563, discloses a process for the simultaneous purification and concentration of the dextran-sucrase enzyme from the culture medium on sucrose. In particular, the disclosed method involves the addition of a polyether such as polyethylene glycol so as to form two phases; the first a heavy dextran-rich phase that contains the concentrated and purified dextran-sucrase enzyme, and the second a lighter polyether-rich phase that contains contaminating enzymatic activities, which is eliminate.

-6-

In view of the above, it is apparent that the cited art does not disclose the recovery and purification of naturally produced chymosin from aqueous solutions containing pepsin and other contaminants by employing an aqueous two phase system derived from the addition of polyethylene glycol and inorganic salt coupled with the use of an ion exchange resin. On the other hand, industrial or commercial scale recovery and purification of naturally produced chymosin is greatly facilitated by using such a liquid-liquid two phase system for the recovery of chymosin and by using an ion exchange resin for the purification of chymosin.

Accordingly, it is an object of this invention to provide efficient processes for the recovery and purification of naturally produced chymosin from aqueous mixtures containing chymosin, pepsin and other contaminants produced by the aqueous extraction of natural chymosin sources.

It is a further object of this invention to provide a recovery and purification process for naturally produced chymosin wherein recovery is achieved by using a liquid-liquid two phase system and purification is achieved by using an ion exchange resin.

It is still a further object of this invention to provide a recovery and

-7-

purification process for naturally produced chymosin wherein recovery is achieved by using a liquid-liquid two phase system which provides for the selective recovery of
5 chymosin and pepsin from other contaminants found in the aqueous solution.

It is a further object of this invention to provide a purification process for naturally occurring chymosin from an aqueous/poly-
10 thylene glycol solution containing chymosin and pepsin.

These and other objects are achieved by the present invention as evidenced by the attached summary of the invention, detailed
15 description of the invention, examples, and claims.

SUMMARY OF THE INVENTION

In one aspect, this invention is a method for recovering and purifying chymosin from an
20 aqueous solution which contains chymosin obtained from natural sources and which additionally contains pepsin and other contaminants which method comprises:

a) adding to the aqueous solution an
25 effective amount of polyethylene glycol (PEG) and an inorganic salt so as to form a two phase system;

b) allowing the aqueous solution-polyethylene glycol-inorganic salt mixture to

-8-

separate into a chymosin and pepsin-rich polyethylene glycol phase and a chymosin and pepsin-poor salt phase;

5 c) recovering the chymosin and pepsin-rich polyethylene glycol phase,

d) contacting the chymosin and pepsin-rich polyethylene glycol phase with an ion exchange resin under conditions wherein the chymosin binds to the resin and the
10 polyethylene glycol and pepsin pass through the resin; and

e) recovering the chymosin from the resin.

15 Surprisingly, in the extraction step, most of the chymosin and pepsin are partitioned into the polyethylene glycol phase whereas the contaminants other than pepsin remain in the salt phase. Moreover, when the
20 polyethylene glycol phase is contacted with a suitable ion exchange resin, the chymosin binds to the resin whereas the pepsin passes through. Afterwards, the chymosin, substantially free of pepsin and other
25 contaminants (i.e, the amount of contamination is less than about 10% by weight based on the weight of chymosin), is recovered from the resin. Thus, the method of the present invention is an extremely efficient recovery
30 and purification method for naturally produced chymosin.

-9-

In general, the pH of the aqueous solution can be any pH at which the chymosin is stable, i.e., about 6.5 or less. However, in a preferred embodiment, it has been found that the use of lower pHs, i.e., pH 3 or less and preferably from about pH 2 to about 2.5, in the aqueous solution result in higher partition coefficients (higher selectivity) for the separation of chymosin into the polyethylene glycol phase as compared to the use of pHs of from above 3 to about 6.5.

Accordingly, a preferred method aspect of the present invention relates to a method for recovering and purifying chymosin from an aqueous solution which contains chymosin obtained from natural sources and which additionally contains pepsin and other contaminants which method comprises:

- a) adjusting the pH of the aqueous solution to less than about 3 and then adding to the aqueous solution an effective amount of polyethylene glycol (PEG) and an inorganic salt so as to form a two phase system;
- b) allowing the aqueous solution-polyethylene glycol-inorganic salt mixture to separate into a chymosin and pepsin-rich polyethylene glycol phase and a chymosin and pepsin-poor salt phase;
- c) recovering the chymosin and pepsin-rich polyethylene glycol phase,
- d) contacting the chymosin and

-10-

pepsin-rich polyethylene glycol phase with a cation exchange resin under conditions wherein the chymosin binds to the resin and the polyethylene glycol and pepsin pass through the resin; and

e) recovering the chymosin from the resin.

DETAILED DESCRIPTION OF THE INVENTION

The separation of various enzymes from aqueous solutions using polymers such as polyethylene glycol of various molecular weights in combination with other polymers, e.g., dextran, or inorganic salts is known in the art. However, this invention is directed in part to the unexpected discovery that chymosin can be efficiently recovered and purified from aqueous solutions containing naturally produced chymosin, pepsin and other contaminants by adding a sufficient amount of both polyethylene glycol and an inorganic salt to the aqueous solution so as to form a two phase system. Under these circumstances, almost all of the chymosin and pepsin is partitioned into the polyethylene glycol phase. This is evidenced by partition coefficients for chymosin and pepsin in the polyethylene glycol phase of greater than about 30 and preferably greater than about 85. On the other hand, most of the other contaminants remain in the salt phase. Thus, this extraction step provides a means of recovering

-11-

the chymosin and separating it from
contaminants other than pepsin.

5 The present invention is further directed
in part to the unexpected discovery that when
the polyethylene glycol phase is then
contacted with an ion exchange resin under
conditions wherein the chymosin binds to the
resin, the polyethylene glycol and pepsin will
pass through the resin. Accordingly, when
10 these two steps are combined, the method of
the present invention permits the recovery and
purification of chymosin which is
substantially free of pepsin and other
contaminants.

15 However, prior to discussing this
invention in detail, the following terms will
first be defined:

20 "Naturally produced chymosin" -- refers
to chymosin obtained from mammalian sources
including, for example, the stomachs of bovine
(including the fourth stomachs of calves), of
goat, of porcine, of lamb, etc.

25 "Pepsin" -- refer to pepsin obtained from
mammalian sources including, for example, the
stomachs of bovine (including the fourth
stomach of calves), of goat, or porcine, of
lamb, etc. In addition to chymosin, pepsin is
also recovered in the extraction of macerated
stomachs.

-12-

"Other contaminants" -- refer to components, other than chymosin and pepsin, obtained during the extraction of mammalian macerated stomachs. Such other contaminants include, for example, proteins (e.g., albumin), enzymes and the like.

"Aqueous solutions containing naturally produced chymosin, pepsin and other contaminants" -- refers to aqueous solutions obtained by extracting macerated mammalian stomach tissue so as to obtain naturally produced chymosin. Methods for preparing (macerating) and extracting chymosin from such tissue are well known in the art and are described, for example, in "Fundamentals of Dairy Chemistry", 2nd Ed., Webb et al. Editors, AVI Publishing Company, pp. 674-679, (1983). Such known extractions procedure also extract pepsin from the stomach tissue as well as other contaminants. In general, the amount of pepsin extracted into the aqueous solution depends on factors such as the age of the animal from which the stomach was obtained as well as the whether the animal had been weaned. The stomach of a young animal which is still nursing will contain significantly more chymosin and less pepsin as compared to an older animal which had been weaned. On the other hand, the stomachs of such older animals still contain recoverable amounts of chymosin.

-13-

"Polyethylene glycol" -- refers to any molecular weight polyethylene glycol which can be used to extract chymosin in the manner of this invention. Polyethylene glycol is
5 available in molecular weights ranging from about 400 to about 22,000. Preferred polyethylene glycol for use herein should have a molecular weight in the range from about 600 to about 12,000. A particularly preferred
10 polyethylene glycol is PEG-8000, i.e., polyethylene glycol having a molecular weight in the range of about 8,000. The selection of the polyethylene glycol used will depend in part on the composition of the mixture from
15 which the chymosin is to be extracted and in part on economics of the process, as well as other factors.

"Inorganic salt" -- refers to any inorganic salt which can be used to extract
20 chymosin in the manner of this invention. Suitable inorganic salts include for instance, sulfate salts, phosphate salts, and the like. The sulfate salts are preferred including sodium sulfate, magnesium sulfate, ammonium
25 sulfate, and the like. Additionally, mixtures of suitable salts can also be used as well as mixtures of such salts in combination with salt(s) such as sodium chloride, which by itself does not partition into a two phase
30 system with polyethylene glycol but in combination with a suitable inorganic salt are

-14-

known to enhance the partition coefficients of enzymes.

"Partition coefficient (K)" -- is defined by the formula

5
$$K = C_t/C_b$$

where C_t refers to the equilibrium concentration of the partitioned compound in the top phase and C_b refers to the equilibrium concentration of the partitioned compound in the bottom phase. Accordingly, it is apparent that the quantitative amount of partitioned compound in either phase depends on its partition coefficient as well as the volume of the phases. That is to say that if the partitioned compound has a partition coefficient of unity (the compound is equally partitioned in the top and bottom phases), then the phases will contain equal quantities of the partitioned compound only if the phases are of equal volume. If the top phase has 10% of the volume of the bottom phase, then when the partition coefficient is unity, the top phase will contain only 10% of the partitioned compound. In view of the above, it is further apparent that a very high partition coefficient for the partitioned compound is extremely valuable because it allows recovery of large quantities of this compound in the upper phase even when the volume of the upper phase is relatively small as compared to the bottom phase. Thus, in the present invention, very high partition coefficients allow for the

10
15
20
25
30

-15-

use of smaller quantities of polyethylene glycol while still achieving very high recoveries of chymosin.

5 "Isoelectric point (IP)" -- refers to the
pH at which a polypeptide will be electro-
statically neutral, i.e., the polypeptide
carries an equal number of positive and
negative charged functionalities. The
isoelectric point for chymosin is about 4.6
10 and that for pepsin is also about 4.6. At a
pH below the isoelectric point, chymosin and
pepsin will have a net positive charge; and at
a pH above the isoelectric point, chymosin and
pepsin will have a net negative charge.

15 "Ion exchange resin" -- refers to a
protein compatible resinous material which is
capable of electrostatically binding charged
compounds. Ion exchange resins are well known
in the art and include both cation and anion
20 exchange resins.

In the practice of this invention, an
aqueous polyethylene glycol solution
containing chymosin and pepsin is contacted
with an ion exchange resin under conditions
25 wherein the chymosin will bind to the resin.
Whether a cation or an anion exchange resin is
employed in the present invention depends on
the pH of the polyethylene glycol phase, i.e.,
whether the pH of the solution is above or
30 below the isoelectric point of chymosin.

-16-

Accordingly, contacting a solution containing chymosin with an ion exchange resin under conditions wherein the chymosin binds to the resin merely refers to adjusting the pH of the solution above or below its isoelectric point so that the chymosin binds to the resin employed.

On the other hand, it has unexpectedly been found that under these conditions, pepsin does not bind to the ion exchange resin even though pepsin has an isoelectric point similar to that of chymosin.

The pH of the aqueous polyethylene glycol solution is generally about 6.5 or less, although pH's around chymosin's isoelectric point, i.e., pH from about 3.6 to about 5.0, are not preferred due to the low net electrostatic charge of chymosin which reduces its effectiveness in binding to the resin. Additionally, when the polyethylene glycol solution (phase) is maintained at a pH between about 3-5, chymosin undergoes more efficient autolysis, although at a rate appreciably slower than in water. In any event, maintaining the aqueous polyethylene glycol phase at a pH between about 3-5 will result in some loss in chymosin yield due to autolysis. Accordingly, when a cation exchange resin is employed, it is preferred that the pH of the solution be maintained below about 3.0; whereas when an anion exchange resin is

-17-

employed, it is preferred that the pH be maintained at above about 5.0.

Preferred cation exchange resins for use in this invention include, for instance, IBF SP-Spherodex, Pharmacia SP-Sephadex, Indion SP-2, IBF SP-Trisacryl, and the like. Preferred anion exchange resins for use in this invention include, for instance, IBF Q-Spherodex, Pharmacia Q-Sephadex, Indion Q-2, IBF Q-Trisacryl, and the like.

The processes of this invention are useful for recovery and purification of naturally produced chymosin. When recovering and purifying naturally produced chymosin, the aqueous solution containing chymosin, pepsin and other contaminants may be used in its crude form, i.e., the solution obtained from extracting macerated stomach, or, if desired, the aqueous solution can first be filtered to remove most or all of the solids then the liquid filtrate used in the processes of this invention.

In the first step of this invention, naturally produced chymosin is recovered by adding to the aqueous solution containing chymosin, pepsin and other impurities, an effective amount of polyethylene glycol (PEG) and an effective amount of an inorganic salt so as to form a two phase system. The resulting solution is allowed to stand so as

-18-

to separate into a chymosin and pepsin-rich polyethylene glycol phase and a chymosin and pepsin-poor salt phase. The chymosin and pepsin rich polyethylene glycol phase is then
5 recovered by conventional techniques.

It has been found that under these conditions, partition coefficients for chymosin and pepsin in the polyethylene glycol phase of greater than 30 and preferably
10 greater than 85 are achieved. While polyethylene glycol extraction is useful for recovering chymosin at higher pH levels, i.e., pH 6.5 or less, it is more efficient as when the process is conducted at a pH less than
15 about 3. At such lower pHs, partition coefficients for chymosin/pepsin of up to 100 or more can be achieved. High partition coefficients are particularly advantageous because such permit the use of smaller
20 quantities of polyethylene glycol to achieve the desired separation of chymosin from the fermentation beer which, in turn, facilitates the later separation of chymosin from the polyethylene glycol, i.e., there is less
25 polyethylene glycol to separate.

It has further been found that as a result of a single extraction, the polyethylene glycol phase can contain as much as 95% or more of the total chymosin initially
30 present in the aqueous solution and contains very little, if any, of the other

-19-

contaminants. Thus, in addition to providing a means for recovering substantially all of the naturally produced chymosin and pepsin contained in the aqueous solution, this aspect
5 of the present invention also provides a means of recovering the chymosin and separating it from contaminants other than pepsin.

In the next step of this invention, the polyethylene glycol phase containing the
10 extracted chymosin and pepsin is separated from the other phase or phases and the polyethylene glycol phase is contacted with an ion exchange resin while maintaining or adjusting the pH so that the chymosin will
15 bind to the resin. Because the polyethylene glycol is not charged, under these conditions, it passes through the resin. Surprisingly, under these conditions, the pepsin also does not bind to the ion exchange resin and passes
20 through the resin. Thus, this step effects purification of chymosin not only from the polyethylene glycol but also from the pepsin. Thus, when the isolated polyethylene glycol phase containing the extracted chymosin and
25 pepsin is contacted with the ion exchange resin under conditions where the chymosin binds to the resin, essentially all the chymosin comes out of the polyethylene glycol and is bound to the ion exchange resin and the
30 polyethylene glycol and pepsin pass through the resin column. After the initial contact, the resin is washed with either water or water

-20-

and salt, preferably under conditions that will not remove the chymosin from the resin, to remove the remaining polyethylene glycol and pepsin. Then the chymosin is eluted from the column using a salt solution and a buffer maintained at a pH which will remove the chymosin from the resin. Because of the high selectivity for chymosin recovery, it is unnecessary to use a gradient or step-wise elution of the resin because essentially the only enzyme or material which is bound by the resin from the polyethylene glycol phase and subsequently released from the resin is chymosin. Therefore, the chymosin can be eluted in one bulk step using the salt solution and raising or lowering the pH (depending of course, on whether a cation or anion exchange resin is employed) to cause the entire chymosin content bound to the resin to be eluted in one batch. Preferably, a salt is added to the eluting solution to aid in the rate or degree of elution, or in some cases, i.e., with an anion exchange resin, to effect elution of the chymosin from the resin, e.g. 50mM sodium phosphate/2M NaCl, pH 5.8. Salts are preferably employed with the eluting solution since chymosin is usually sold in commercial form in a salt solution and accordingly, it is convenient to incorporate the salt with the chymosin at this point.

Thus, recovery and purification of chymosin in the manner of this invention will result in chymosin substantially free of

-21-

pepsin and other contaminants, i.e., the amount of contamination is less than about 10% by weight based on the weight of chymosin and preferably, less than about 5% by weight based on the weight of the chymosin. That is to say that the resulting chymosin product is at least about 90% by weight pure and can be prepared for commercial use without further significant treatment to remove impurities.

The commercial chymosin product is usually diluted to about 5 grams per gallon or about 1.5 grams per liter chymosin. The salt (usually NaCl) concentration is normally brought up to about 18% and a preservative such as sodium benzoate is added. The final concentration product intended for food grade use usually is also subjected to a final filtration to remove and undesirable solids or particles that may be present.

Although higher pH levels can be used throughout the process (pHs up to about 6.5), the efficiency of the polyethylene glycol/inorganic salt mixture in extracting the chymosin from the aqueous solution containing chymosin, pepsin and other contaminants and therefore the efficiency of the process is not as high as when low pH is maintained throughout the extraction step. Therefore, it is preferred to use low pH and accordingly, a cation exchange resin to increase the efficiency of the overall process.

-22-

When the preferred aspects of the above invention are combined by conducting the polyethylene glycol extraction at a pH of about 3 or below and contacting the separated polyethylene glycol phase with a cation exchange resin while maintaining the low pH, it has been found that a single polyethylene glycol extraction and a single-pass contact with the ion exchange resin will recover as much as 90 to 95% of the total chymosin present in the initial aqueous solution.

On the other hand, the use of a higher pH in the polyethylene glycol extraction step as well as an anion exchange resin while maintaining the pH above chymosin's isoelectric point also provides acceptable results. However, if a higher pH is employed to extract the chymosin and a lower pH is desired to contact the solution with resin, this can be readily achieved simply by adjusting the pH of the polyethylene glycol extract to below chymosin's isoelectric point and preferably, to about pH 3.6 or below and more preferably about pH 3 or below, and then contacting the chymosin with a cation exchange resin.

In addition to the high efficiency for chymosin and pepsin, especially chymosin, recovery by using the polyethylene glycol/inorganic salt mixture herein

-23-

described, it has been further found that the solubility of chymosin in polyethylene glycol is apparently so high that the chymosin moves from the aqueous phase into the polyethylene glycol phase very rapidly. The time required for extraction of the chymosin into the polyethylene glycol phase is usually so short that it is not a significant process design factor. This process is therefore very efficient and economic in operation and is easily scaled up for commercial production.

As is apparent and regardless of the pH employed, the herein described process involves moving the chymosin and pepsin from the aqueous mixture into the more hydrophobic polyethylene glycol phase. This is driven, at least in part, by the salt concentration in the non-polyethylene glycol phase or phases. If a low molecular weight polyethylene glycol is used, the polyethylene glycol phase is less hydrophobic and a higher salt concentration is necessary in the non-polyethylene glycol phase(s), which adds to the cost of operation. If a higher molecular weight more hydrophobic polyethylene glycol is used, less salt will be needed in the process, but the separation rate may be lower because of the high viscosity of the higher molecular weight polyethylene glycol. Thus, the process of this invention can be optimized for any particular operation by selecting the desired polyethylene glycol molecular weight, the salt concentration and

-24-

other parameters which provide the desired economics. One objective is usually to minimize the time required to move the chymosin into the polyethylene glycol phase, but another objective is usually to minimize the amount of salt used to effect the transfer of essentially all of the chymosin into the polyethylene glycol phase. While the salt concentration in the non-polyethylene glycol phases can be 20% by weight or higher, usually less than about 15% is required with the appropriate polyethylene glycol. For example, with PEG-8000, about 10-13% sodium sulfate is adequate. On the other hand, the minimum inorganic salt concentration is dictated by the concentration of the salt necessary to form a two phase system with polyethylene glycol. However, in a preferred embodiment, the inorganic salt concentration is from about 8.5 to about 20 weight to volume percent based on the volume of the aqueous solution containing chymosin, pepsin and other contaminants.

Also, preferably, the polyethylene glycol concentration employed herein is less than about 20, and more preferably less than about 15, weight to volume percent based on the volume of the aqueous solution containing chymosin, pepsin and other contaminants. For economics and ease of later separation, as little polyethylene glycol as possible is most preferably employed.

-25-

The exact concentration of polyethylene glycol and inorganic salt employed herein can readily be determined by the skilled artisan.

5 After the chymosin is bound by the ion exchange resin, the recovered polyethylene glycol phase is contaminated with pepsin. The pepsin can be recovered from this solution by known methods and the polyethylene glycol recycled. Alternatively, the polyethylene glycol can be discarded.

10 On the other hand, the ion exchange resin can be regenerated for use with subsequent batches of polyethylene glycol containing chymosin by washing the ion exchange resin with a solution of water adjusted to the appropriate pH. For example, when a cation exchange resin is employed, it can be regenerated by washing it with a solution of water containing enough sulfuric acid to make the pH about 2.

15 The above aspects of this invention which enable the reuse of the ion exchange resins particularly lend the processes of this invention to efficient commercial and industrial operation for the purification of industrial quantities of chymosin, particularly naturally produced chymosin.

-26-

Having described the invention in general terms the invention can be better understood by reference to the following embodiments of the invention which are illustrated in the following examples. However the scope of this invention is to be determined by the appended claims, whereas the following examples are merely illustrative embodiments of particular ways in which the invention disclosed herein can be practiced.

EXAMPLES

Example 1

This example describes the use of an aqueous two phase polyethylene glycol extraction process followed by contact with an ion exchange resin to produce food grade chymosin from bovine stomachs. The chymosin is recovered from beef (bovine) stomachs which contain large amounts of bovine pepsin and a lesser amount of chymosin. The aqueous solution employed herein is prepared by grinding bovine stomachs in a suitable liquid medium and filtering out the cell debris. The resulting aqueous solution contains naturally produced chymosin, pepsin and other contaminants.

In this example, 500 ml of beef extract was adjusted to pH 2 with sulfuric acid. 20 grams of PEG 8000 (4 wt. to vol. percent) and

-27-

55 grams of anhydrous sodium sulfate (11 wt. to vol. percent) were added to the extract. The extract was warmed to 37°C to facilitate the solubility of the sulfate salt. The mixture was separated into two phases by centrifugation (Sorvall centrifuge) at about 5000 x g for about 15 minutes and the polyethylene glycol phase (top phase) was separated from the salt phase (bottom phase) by removing the bottom phase with a peristaltic pump. The polyethylene glycol phase was diluted 1:3 with deionized water. The pH of this phase was determined to be 2.7 and was adjusted to 2.3 with sulfuric acid. The polyethylene glycol diluted phase was then passed over an ion exchange resin consisting of IBF Spherox SP (a protein compatible cation exchange resin) which was previously equilibrated in water at pH 2 so as to bind the chymosin. The polyethylene glycol solution which flowed through was collected and was found to contain pepsin. The resin was eluted with a solution of 0.5 M NaCl at pH 2.0 to remove any remaining polyethylene glycol and pepsin and the wash solution was collected. The resin was then eluted with a solution of 0.05 M sodium phosphate at pH 5.8 containing 2 M NaCl so as to release the chymosin in bulk. The eluted liquid containing chymosin substantially free of pepsin and other contaminants was then recovered.

-28-

For the purpose of distinguishing the relative amounts of bovine pepsin and chymosin in the product, the following assay was employed. The milk clotting ability of the various solutions was measured using a roller bottle apparatus. Skim milk was prepared at pH 6 and at 6.5. Bovine pepsin has more activity at pH 6 and chymosin has more activity at pH 6.5. The clotting activities are compared to a standard consisting of 90% chymosin, 10% pepsin. The ratio of activity at pH 6 to pH 6.5 is then a measure of the relative amounts of the two enzymes. The standard, by definition, provides a ratio of 1. Since bovine pepsin is more active at pH 6 than at pH 6.5, the ratio of activity at pH 6 and pH 6.5 will be greater than 1 and approach 2 for pure pepsin. On the other hand, for high levels of chymosin, the ratio will be less than 1 and approach 0.5 or less.

In this assay, the aqueous solution of chymosin, pepsin and other impurities had a total concentration of chymosin and pepsin of 16.64 CHU/ml (pH 6.5). After recovering the top phase, it was applied to the resin. The results of the resin wash are reported as flow through (the material which passed through the resin without binding); wash (the material which came off the resin during the washing step to remove pepsin and other contaminants); and the elute (the material recovered when the

-29-

column was eluted so as to recover chymosin).
The results of this assay are as follows:

-30-

	<u>Sample</u> <u>assayed</u>	<u>Conc. of chymosin</u> <u>and pepsin</u>	<u>ratio (6/6.5)</u>
5	starting material	-----	1.41
	extraction of 500 ml	8320 CHU ^a (16.54 CHU/ml)	
	top phase ^b	7830 CHU (115.15 CHU/ml)	1.34
	bottom phase	454.5 CHU (1.01 CHU/ml)	---
10	flow through	971.6 CHU (6.94 CHU/ml)	1.08
	wash	179.75 CHU (7.19 CHU/ml)	1.54
	elute	822.25 CHU (32.89 CHU/ml)	0.49

a = CHU -- Chris. Hansen Unit - 1 CHU/ml
equals under the following conditions:

15 Substrate: 110 g of low heat, spraydried
skim milk powder is suspended in 1000
ml 0.05% calcium chloride. The milk is
20 stirred for 30 minutes at room
temperature and then left for rest for
another 30 minutes. The milk should be
stored at a temperature between 4 and
25°C and not longer than 3 hours. The
pH of the milk is about 6.5.

25 Temperature: 32°C plus or minus 0.2°C in
a thermostatic water bath.

Enzyme Addition: To 25 ml of the
reconstituted skim milk is added 0.5 ml
of enzyme solution, diluted to give a
clotting time of between 380 and 500
30 seconds,

will give a clotting time of 410 to 460
seconds.

b = the partition coefficient for
chymosin/pepsin is 114

-31-

From the above data, it is apparent that the material recovered in the elute is chymosin substantially free of pepsin and other contaminant. Thus, the process of this invention permits the efficient recovery of chymosin from natural chymosin sources.

Likewise, by following the procedures set forth above, chymosin can likewise be recovered and purified by extracting at a pH of from about 5 to 6.5 and using an anion exchange resin. Suitable anion exchange resins which could be substituted for the cation exchange resin in the above example include, for instance, IBF Q-Spherox, Pharmacia Q-Sephadex, Indion Q-2, IBF Q-Trisacryl, and the like.

Example 2

In a manner similar to the extraction procedure set forth in Example 1 above but using a pH of about 5.8, the enzymes listed below were separately extracted from an aqueous solution into the polyethylene glycol phase of the liquid-liquid two phase system to provide the following results (pure enzyme used):

-32-

	Enzyme	Protein (mg/ml)	Activity CHU/ml	Partition Coefficient
	calf chymosin	1.7	28	87
	bovine pepsin	1.9	31	31.5
5	porcine pepsin	2.3	29.7	1.45
	E. parasitica aspartic protease	3.6	27	0.33
10	M. miehei aspartic protease	2.4	20.5	0.24

This experiment was repeated but at this time at pH 2-2.5 with the following results:

	Enzyme	Protein (mg/ml)	Activity CHU/ml	Partition Coefficient
15	calf chymosin	1.2	89	>943 ^c
	bovine pepsin	1.3	50	>462 ^c
	porcine pepsin	1.6	23.7	>206 ^c
20	E. parasitica aspartic protease	2.5	15.2	1.18
	M. miehei aspartic protease	1.7	48	0.94

25 c = Partition coefficients of about 200 or more are difficult to measure accurately because of assay limitations. That is to say that because the partition coefficient is a ratio of the concentration of chymosin in the top phase divided by the concentration of the chymosin in the bottom phase and further because the amount of chymosin in the bottom phase is generally very small, small changes in this bottom concentration will produce large swings in the partition coefficient. Moreover, the concentration determined by assay methodology is particularly subject to variations at very low concentrations.

30

35

-33-

The above results indicate that chymosin and pepsin have high partition coefficients in this system. However, it is also clear that the partition coefficients for bovine pepsin increase substantially in going from pH 5.8 to pH 2-2.5. Accordingly, it may be preferably to conduct the extraction at a pH above about 5 so as to reduce the amount of pepsin in the polyethylene glycol phase. On the other hand, the partition coefficients for chymosin also increase substantially in going from pH 5.8 to pH 2-2.5. Thus, if the extraction is done at a pH above about 5, the amount of chymosin recovered would also somewhat reduced, albeit at a lower level because chymosin's partition coefficient at pH 5.8 is higher than that of bovine pepsin.

-34-

WHAT IS CLAIMED IS:

- 1 1. A method for recovering and
2 purifying chymosin from an aqueous solution
3 which contains chymosin obtained from natural
4 sources and which additionally contains pepsin
5 and other contaminants which method comprises:
6 a) adding to the aqueous solution an
7 effective amount of polyethylene glycol (PEG)
8 and an inorganic salt so as to form a two
9 phase system;
10 b) allowing the aqueous solution-
11 polyethylene glycol-inorganic salt mixture to
12 separate into a chymosin and pepsin-rich
13 polyethylene glycol phase and a chymosin and
14 pepsin-poor salt phase;
15 c) recovering the chymosin and
16 pepsin-rich polyethylene glycol phase,
17 d) contacting the chymosin and
18 pepsin-rich polyethylene glycol phase with an
19 ion exchange resin under conditions wherein
20 the chymosin binds to the resin and the
21 polyethylene glycol and pepsin pass through
22 the resin; and
23 e) recovering the chymosin from the
24 resin.
- 1 2. A method according to Claim 1
2 wherein the pH of the aqueous solution is
3 about 6.5 or less.

-35-

1 3. A method according to Claim 2
2 wherein the pH of the aqueous solution is
3 about 3 or less.

1 4. A method according to Claim 3
2 wherein the pH of the aqueous solution is less
3 than about 2.8.

1 5. A method according to Claim 1
2 wherein the average molecular weight of the
3 polyethylene glycol is from about 600 to about
4 12,000.

1 6. A method according to Claim 5
2 wherein the average molecular weight of the
3 polyethylene glycol is from about 5,000 to
4 about 10,000.

1 7. A method according to Claim 1
2 wherein said inorganic salt is selected from
3 the group consisting of sulfate salts, and
4 phosphate salts.

1 8. A method according to Claim 7
2 wherein said inorganic salt is a sulfate salt.

1 9. A method according to Claim 8
2 wherein said sulfate salt is selected from the
3 group consisting of sodium sulfate, magnesium
4 sulfate, and ammonium sulfate.

1 10. A method according to Claim 1
2 wherein said aqueous solution is first

-36-

3 filtered prior to addition of said
4 polyethylene glycol and said inorganic salt.

1 11. A method according to Claim 1
2 wherein the pH of said chymosin and pepsin-
3 rich polyethylene glycol phase is either from
4 5.0 to 6.5 or is about 3.0 or less.

1 12. A method according to Claim 11
2 wherein the pH of said chymosin and pepsin-
3 rich polyethylene glycol phase is about 3.0 or
4 less and a cation exchange resin is employed.

1 13. A method according to Claim 12
2 wherein the pH of said chymosin and pepsin-
3 rich polyethylene glycol phase is from about
4 2.0 to 2.5.

1 14. A method according to Claim 11
2 wherein the pH of said chymosin and pepsin-
3 rich polyethylene glycol phase is from about
4 5.0 to about 6.5 and an anion exchange resin
5 is employed.

1 15. A method according to Claim 1
2 wherein the extraction step a) is conducted at
3 a pH of about 3.0 or less and after isolation
4 of the polyethylene glycol phase, the pH of
5 this phase is adjusted to from about 5.0 to
6 about 6.5 and an anion exchange resin is
7 employed.

-37-

- 1 16. A method for recovering and
2 purifying chymosin from an aqueous solution
3 which contains chymosin obtained from natural
4 sources and which additionally contains pepsin
5 and other contaminants which method comprises:
6 a) adjusting the pH of the aqueous
7 solution to less than about 3 and then adding
8 to the aqueous solution an effective amount of
9 polyethylene glycol (PEG) and an inorganic
10 salt so as to form a two phase system;
11 b) allowing the aqueous solution-
12 polyethylene glycol-inorganic salt mixture to
13 separate into a chymosin and pepsin-rich
14 polyethylene glycol phase and a chymosin and
15 pepsin-poor salt phase;
16 c) recovering the chymosin and
17 pepsin-rich polyethylene glycol phase,
18 d) contacting the chymosin and
19 pepsin-rich polyethylene glycol phase with a
20 cation exchange resin under conditions wherein
21 the chymosin binds to the resin and the
22 polyethylene glycol and pepsin pass through
23 the resin; and
24 e) recovering the chymosin from the
25 resin.

- 1 17. A method according to Claim 16
2 wherein the average molecular weight of the
3 polyethylene glycol is from about 600 to about
4 12,000.

- 1 18. A method according to Claim 17
2 wherein the average molecular weight of the

-38-

3 polyethylene glycol is from about 5,000 to
4 about 10,000.

1 19. A method according to Claim 16
2 wherein said inorganic salt is selected from
3 the group consisting of sulfate salts, and
4 phosphate salts.

1 20. A method according to Claim 19
2 wherein said inorganic salt is a sulfate salt.

1 21. A method according to Claim 20
2 wherein said sulfate salt is selected from the
3 group consisting of sodium sulfate, magnesium
4 sulfate, and ammonium sulfate.

1 22. A method according to Claim 16
2 wherein said aqueous solution beer is first
3 filtered prior to addition of said
4 polyethylene glycol and said inorganic salt.

INTERNATIONAL SEARCH REPORT

International Application No PCT/US90/03378

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all)³

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(5): C12N 9/64

U.S.Cl.: 435/226,815,816

II. FIELDS SEARCHED

Minimum Documentation Searched⁴

Classification System:

Classification Symbols

U.S. 435/226,815,816
426/36,42

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched⁴

COMPUTER SEARCH-APS, CA, DIALOG

III. DOCUMENTS CONSIDERED TO BE RELEVANT¹⁴

Category ⁵	Citation of Document, ¹⁵ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁶
Y	US, A, 4,144,130 (Kula, et. al.) 13 March 1979. See entire document.	1,2,5-10,14,15
Y	US, A, 4,728,613 (Brewer et. al.) 01 March 1988. See especially column 2, line 50. See entire document.	1,2,5-10,14,15
A	US, A, 4,743,551 (Subramanian) 10 May 1988.	1-22
A	US, A, 4,745,063 (Birschbach) 17 May 1988.	1-22

⁵ Special categories of cited documents: ¹⁵

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search¹

05 September 1990

Date of Mailing of this International Search Report¹

22 OCT 1990

International Searching Authority¹

ISA/US

Signature of Authorized Officer¹⁰

Charles Patterson
Charles Patterson